

Embracing IGRA's in Children

Deborah Lewinsohn, MD

Associate Professor

Infectious Disease, Pediatrics



OREGON
HEALTH & SCIENCE
UNIVERSITY

Global Burden of Tuberculosis

Persons Infected

2 Billion

Estimated Cases 2003


Adults: 8,800,000

Children: 250,000

Estimated Deaths 2003


Adults: 1,700,000

Children: 100,000



WHO: The World Health Report 2007

Natural History of *M. tuberculosis* Infection



Diagnosis of LTBI using TST



*Photographs: Dr. Chuck Daley,
National Jewish Medical Center*

Problems include:


- Inter-reader variability; requires return visit.
- Confusing cutoffs for various risk groups.
- Low positive predictive value in countries with low prevalence
- False negatives
 - Anergy: HIV/ESRD
 - Recent TB
 - Very young
 - Overwhelming disease
- False positives
 - BCG vaccination
 - Nontuberculous mycobacteria

T cell based diagnostics for TB: IFN- γ Release Assays (IGRA's)

- Potent pro-inflammatory cytokine released by T cells and NK cells. Assays reflect adaptive T cell response to TB.
- Two commercially available tests:
 - T-spot[®].TB; ELISPOT.
 - QuantiFERON[®]-TB Gold IT; ELISA.

Antigens Absent from BCG

BCG Deletion (RD01)



* Mtb11 (38.1)
CFP-10
ESAT-6

Gene Map from Sanger Centre; Cole et al., Nature 393, 537-544 (1998).

TST vs IGRA


	TST	IFN
Cell Types	CD4 CD8 Basophils DC and Macrophages	CD4 (CD8)
Cytokines	IL-4, IFN- γ TNF- α , IL-10, IL-12, G-CSF,	IFN- γ
Associated with Protective Immunity	No	Yes
Timing	2-5 Days	Short Term
Homing Phenotype	Skin	Polymorphic

Epithelium or Alveolus


Lymph Node

T Cell Priming

Mtb



Danger



IL-12

TGF-β1
IL-23
IL-6

IL-10

IL-4

Effector T Cells



IFN-γ
TNFα



IFN-γ
TNFα
IL-2

Protective Immunity



IL-17
TNF-α
GM-CSF




IL-10



IL-4
IL-5
IL-13

Non-Protective Immunity

QuantiFERON-TB Gold (2nd)




7ml whole blood

ESAT-6

CFP10

Mitogen

QuantiFERON®-TB Gold IT (3rd)



Adapted from QuantiFERON®-TB Gold IT product insert

QuantiFERON®-TB Gold IT: ELISA

Interpretation of test result

Nil [IU/mL]	TB Antigen minus Nil [IU/mL]	Mitogen minus Nil [IU/mL] ¹	QuantiFERON®-TB [IU/mL]	Report/Interpretation
≤ 8.0	< 0.35	≥ 0.5	Negative	<i>M. tuberculosis</i> infection NOT likely
	≥ 0.35 and < 25% of Nil value	≥ 0.5		
	≥ 0.35 and ≥ 25% of Nil value	Any	Positive²	<i>M. tuberculosis</i> infection likely
	< 0.35	< 0.5	Indeterminate³	Results are indeterminate for TB-Antigen responsiveness
≥ 0.35 and < 25% of Nil value	< 0.5			
> 8.0	Any	Any		

From: QuantiFERON®-TB Gold IT product insert

T-spot[®].TB: ELISPOT




**CPT
tubes**

**Transport <
8 hr at room
temperature**



ELISPOT assay



- Adults and children 10 years old and over: one 8mL or two 4mL tubes
- Children 2-9 years old: one 4mL tube
- Children up to 2 years old: 2mL paediatric tube

From: T-spot[®].TB Visual procedure guide

T-spot[®].TB: ELISPOT

Interpretation of test result

- The test result is Positive if (Panel A-Nil) and/or (Panel B-Nil) ≥ 8 spots.
- The test result is Negative if both (Panel A-Nil) and (Panel B-Nil) ≤ 4 spots.
- The test result is Borderline if the highest of the Panel A or Panel B spot count (minus Nil) is 5, 6 or 7.
- The test result is Invalid if:
 - Nil > 10 spots.
 - Mitogen < 20 spots AND both (Panel A-Nil) and (Panel B-Nil) ≤ 4 spots.

From: Summarized from T-spot[®].TB Product insert

Use of IGRA's in adults

- CDC Recommendations: “QFT-G can be used in all circumstances in which the TST is used, including contact investigations, evaluation of recent immigrants who have had BCG vaccination, and TB screening of health-care workers and others undergoing serial investigation for *Mycobacterium tuberculosis* infection. QFT-G usually can be used in place of (and not in addition to the TST)” (MMWR, Dec. 16, 2005, Vol.54.)
- FDA Approvals:
 - QFT TB (11/28/01); QFT TB Gold (12/2/04); QFT TB Gold-IT (10/10/07)
 - T-SPOT.TB (7/25/08)

Evidence-based evaluation of IGRA's vs TST

Evaluation without reference to a gold standard diagnostic for LTBI:

- Sensitivity in culture positive TB.
- Relationship with exposure risk factors.
- Specificity in low risk populations.
- Prospective prognostic studies.

Evaluation vs TST in children: Promise of IGRA's

Evaluation without reference to a gold standard diagnostic for LTBI:

- **Sensitivity in culture positive TB.**
- Relationship with exposure risk factors.
- Specificity in low risk populations.
- Prospective prognostic studies.

QFT-Gold for Diagnosis of LTBI in active TB

QFT version	Study, year, country	Age range	Young/total	Sensitivity of TST in active TB n/N (%)	Sensitivity of IGRA in active TB n/N (%)
QFT Gold	Connell, 2006, Australia	0 - 18	NR/101	9/9 (100)	9/9 (100)
QFT Gold	Okada, 2007, Cambodia	0 - 5	210/210 < 6 yrs	15/19 (79)	10/19 (53)
QFT Gold IT	Connell, 2008, Australia	1 - 19	NR/100	NR	8/9 (89)
QFT Gold IT	Detjen, 2007, Germany	0 - 16	54% < 3 yrs; n = 73	28 /28 (100)	26/28 (93)
QFT Gold IT	Dogra, 2007, India	1 - 12	42/105 (40%) < 5 yrs	5/8 (52) –Cx+ 9/11 (82)-Rx	5/8 (52) – Cx+ 9/11 (82) - Rx
QFT Gold IT	Dominguez, 2007, Spain	0 - 18	15/134 (11%) < 5 yrs	9/9 (100)	6/9 (67)
QFT Gold IT	Kampmann, 2009, UK	2 - 16	26/91 < 5 years	20/24 (83) Cx+	20/25 (80) Cx+

T-spot[®].TB for Diagnosis of LTBI in active TB

Study, year, country	Age range	Young/total	Sensitivity of TST in active TB n/N (%)	Sensitivity of IGRA in active TB n/N (%)
Connell, 2008, Australia	1 - 19	NR/101	NR	9/9 (89)
Detjen, 2007, Germany	0 - 16	54% < 3 yrs; n = 73	28 /28 (100)	26/28 (93)
Dominguez, 2007, Spain	0 - 18	15/134 (11%) < 5 yrs	9/9 (100)	6/9 (67)
Kampmann, 2009, UK	2 - 16	26/91 < 5 years	20/24 (83)	14/24 (58)
Nicol, 2009, S. Africa	0 - NR	204/214 < 3 years	30/58 (52)	23/58 (40)

Evaluation vs TST in children: Promise of IGRA's

Evaluation without reference to a gold standard diagnostic for LTBI:

- Sensitivity in culture positive TB.
- **Relationship with exposure risk factors.**
- Specificity in low risk populations.
- Prospective prognostic studies.

QFT-Gold (IT) for LTBI diagnosis

QFT version	Study, year, country	Age range	Young/total	TST positivity rate n/N (%)	IGRA positivity Concordance TST/IGRA
QFT Gold	Bergamini, 2009, Italy	0-19	39/181, < 4 yrs	58/181 (32)	34/181 (19), K = 0.43
QFT Gold	Connell, 2006, Australia	0 - 18	NR/101	42/42 (100)	11/42 (26); K = 0.30; TST+/QFT-
QFT Gold	Hesseling, 2008, S. Africa 	0 - 5	29/29	15/28 (54)	NR (88.9); K = 0.78; TST+/QFT-
QFT Gold	Mandalakas, 2008, S. Africa	NR, X = 4.4	All HIV+; NR/23	6/23 (26)	NR (75) K = .44; TST+/QFT-
QFT Gold	Okada, 2007, Cambodia 	0 - 5	210/210 : < 6 yrs	47/195 (24)	171/195 (88); K = 0.62; TST+/QFT-
QFT Gold IT	Bergamini, 2009, Italy	0-19	39/181, <4 yrs	74/315 (24)	54/315 (17); K = 0.35
QFT Gold IT	Chun, 2008, Korea 	0-13	NR: Med = 1.7 yr	26/42 (62)	24/42 (57); K = 0.19
QFT Gold IT	Connell, 2008, Australia	1 - 19	NR/100	47/97 (48)	NR; (75); K = .5
QFT Gold IT	Lighter, 2008, USA 	0 - 18	67/207, < 5 yrs	116/204 (56)	112/207 (55); K = .17 TST+/QFT-
QFT Gold IT	Nakaoka, 2006, Nigeria	0 - 14	NR/207	57/206 (28)	49/66 (74); K = .74; TST-/QFT+
QFT Gold IT	Dogra, 2007, India	1 - 12	42/105 , < 5 yrs	10/105 (9.5)	100/105; (95); K = .73
QFT Gold IT	Dominguez, 2007, Spain	0 - 18	15/134 (11%)	115/134 (86)	BCG-:23/40 (58), K=.24
QFT Gold IT	Kampmann, 2009, UK	0-16	56/118; < 5 yrs	57/114 (50)	NR (77); K = .53
QFT Gold IT	Tsiouris, 2006, South Africa	5-15	NA/NR	80/184 (44)	145/184 (79); K=.56; TST+/QFT-

Diagnosis of LTBI/household contacts: QFT:IT





Figure 2. Proportion of children with positive tuberculin skin test (TST) (>10 mm) and QuantiFERON Gold in Tube (QFT-IT) test results, by adult smear positivity. Error bars show 95% confidence intervals.

T-spot[®].TB for LTBI diagnosis


Study, year, country	Age range	Young/total	TST positivity rate n/N (%)	IGRA positivity Concordance TST/QFT
Bergamini, 2009, Italy	0 - 19	57/154 (37)	28/154 (18)	22/154 (14), K= 0.52
Connell, 2008, Australia	1 - 19	NR/100	47/97 (48)	75% K = .51
Dominguez, 2007, Spain	0 - 18	15/134 (11%)	115/134 (86)	BCG-:25/40 (62), K=.33
Hansted, 2009, Lithuania	10 – 17	0/45	27/45 (60)	8/45 (53)
Hesseling, 2008 South Africa 	0-5	29/29	15/28 (54)	NR (46), K=-0.15 TST-/IGRA+
Kampmann, 2009, UK	0-16	56/118; < 5 yrs	57/114 (50)	NR (75); K = .49
Mandalakas, 2008, S. Africa	NR, X = 4.4	All HIV+; NR/23	6/23 (26)	NR (67) K = .33; TST-/IGRA+

Evaluation vs TST in children: Promise of IGRA's

Evaluation without reference to a gold standard diagnostic for LTBI:

- Sensitivity in culture positive TB.
- Relationship with exposure risk factors.
- **Specificity in low risk populations.**
- Prospective prognostic studies.

Specificity of Diagnosis of LTBI in TB suspects: QFT-IT/ T-spot[®].TB vs TST



Specificity of Diagnosis of LTBI in TB suspects: QFT-IT/ T-spot[®].TB vs TST

Table 3. Indicators of diagnostic accuracy for the tuberculin skin test (TST), QFT-IT, T-SPOT, and the combination of QFT-IT and T-SPOT.

Test	Sensitivity, % (95% CI)	Specificity, % (95% CI)	Positive LR (95% CI)	Negative LR (95% CI)	Positive PV, % (95% CI)	Negative PV, % (95% CI)
TST	100 (88–100)	58 (42–73)	2.4 (1.6–3.4)	0.1 ^a (0.01–0.4)	62 (47–76)	100 (85–100)
QFT-IT	93 (77–99)	100 (91–100)	37.1 ^b (5.3–258.0)	0.1 (0.02–0.3)	100 (87–100)	95 (84–99)
T-SPOT	93 (77–99)	98 (87–100)	37.1 (5.3–258.0)	0.1 (0.02–0.3)	96 (81–100)	95 (83–99)
QFT-IT and T-SPOT (both positive)	89 (72–98)	100 (91–100)	35.7 ^b (5.1–248.4)	0.1 (0.04–0.3)	100 (86–100)	93 (81–99)
QFT-IT and T-SPOT (1 positive)	96 (82–100)	98 (87–100)	38.6 (5.6–267.5)	0.04 (0.01–0.3)	96 (82–100)	98 (87–100)
Stepwise approach ^c	93 (77–99)	100 (91–100)	37.1 ^b (5.3–258.0)	0.1 (0.02–0.3)	100 (87–100)	95 (84–99)

NOTE. Calculations were based on 68 children, including 28 children with confirmed tuberculosis (group A) and 40 unvaccinated children without tuberculosis (19 children with confirmed nontuberculous mycobacterial lymphadenitis [group B] and 21 children with other respiratory tract infections [group C]). LR, likelihood ratio; PV, predictive value; QFT-IT, QuantiFERON-TB Gold In-Tube (Cellestis); T-SPOT, T SPOT-TB (Oxford Immunotec).

^a For calculation of this LR, we classified 1 correct diagnosis in a child with tuberculosis as a false-negative result (to avoid division by 0).

^b For calculation of this LR, we classified 1 correct diagnosis in a child without tuberculosis as a false-positive result (to avoid division by 0).

^c We also calculated diagnostic accuracy in a stepwise approach according to recent National Institute of Health and Clinical Excellence guidelines [15]; only patients with positive TST results were tested with IFN- γ release assays. QFT-IT and T-SPOT showed the same values for all indicators.

Specificity of IGRA in BCG-vaccinated children

- Inclusion: No risk and TST ≥ 5 mm.
- 65 children age 2 months – 14 years, all BCG vaccinated.
- 1/65 QFT-IT positive

Evaluation vs TST in children: Promise of IGRA's

Evaluation without reference to a gold standard diagnostic for LTBI:

- Sensitivity in culture positive TB.
- Relationship with exposure risk factors.
- Specificity in low risk populations.
- **Prospective prognostic studies.**

Diagnosis of LTBI/school contacts: QFT

- 349 15-16 y/o boys, all BCG vaccinated.
- Tested with TST: 95 of 349 positive.
- 88 TST positive tested with QFT-TB Gold: 4 of 88 positive.
- 3 of 4 in high exposure group – received INH.
- Remaining TST positive students – no INH and no disease with 3+ years follow-up.


Limitations of IGRA's in Children

- Paucity of data in children < 5 years.
- Increased frequency of indeterminate assays in children < 5 years.
- Required blood volumes.
- Lack of longitudinal data.

Limitations of IGRA's in young children

QFT version	Study, year, country	Age range	Young/total	Indeterminate n/N (%)	QNS Blood n/N (%)
QFT Gold	Connell, 2006, Australia	0 - 18	NR/101	17/101 (17)	3/101 (3)
QFT Gold	Hesseling, 2008, S. Africa	0 - 5	29/29	3/21 (14)	7/29 (24)
QFT Gold	Mandalakas, 2008, S. Africa	NR, X = 4.4	All HIV+; NR/23	0/12 (0)	11/23 (47)
QFT Gold	Okada, 2007, Cambodia	0 - 5	210/210 : < 6 yrs	9/208 (6)	13/217 (6)
QFT Gold IT	Chun, 2008, Korea	0-13	NR: Med = 1.7 yr	17/227 (7.5)	NR
QFT Gold IT	Connell, 2008, Australia	1 - 19	NR/100	0/38 (0)	5/101 (5)
QFT Gold IT	Lighter, 2008, USA	0 - 18	67/207, < 5 yrs	3/207 (1)	0/207 (0)
QFT Gold IT	Nakaoka, 2006, Nigeria	0 - 14	NR/207	33/207 (16)	NR
QFT Gold IT	Dogra, 2007, India	1 - 12	42/105 , < 5 yrs	0/105 (0)	0/105 (0)
QFT Gold IT	Dominguez, 2007, Spain	0 - 18	15/134 (11%)	3/134 (2)	NR
QFT Gold IT	Kampmann, 2009, UK	0-16	56/118; < 5 yrs	14/209 (7)	0/209 (0)
QFT Gold IT	Tsiouris, 2006, South Africa	5-15	NA/NR	NR	37/221 (17)


Indeterminate rates of IGRA's




QFT-TB Gold vs QFT-IT vs T-spot,
Italy

Bergamini, et al., Pediatrics, 2009.

Longitudinal IGRA data in children



Longitudinal IGRA data in children



QFT vs T-SPOT.*TB* in children

- QFT more available than T-SPOT.*TB*
- More published data in children for QFT than for T-SPOT.*TB*.
- Less blood required for T-SPOT.*TB* than for QFT.
- Specificity equivalent in one study.
- Sensitivity of T-SPOT.*TB* increased, equal, decreased when compared with QFT-IT.

Indications for IGRA's and TST in children (My opinion)

- Close contacts of active TB cases:
- Immigrants from endemic countries:
- Significant travel history:
 - IGRA preferred to TST in children ≥ 5 years.
 - TST preferred to IGRA in children < 5 years.

Indications for IGRA's and TST in children (My opinion - continued)

- TB suspects:
- HIV infection:
- Increased risk of progression of LTBI:
 - Consider both IGRA and TST and take either positive as evidence of infection.

Upcoming recommendations

- ATS/CDC/IDSA (AAP) guidelines:
- CDC guidelines – updated for QFT-Gold-IT and T-SPOT.*TB*.
- AAP – 2009 RedBook

Future research needs for IGRA's in children

- High risk young children in low incidence setting (Household contact study in young children).
- Longitudinal data in young children (Data to inform “window prophylaxis”).
- Young immunosuppressed and HIV-infected children.

New assays for TB diagnosis in children

- IP-10 assays (ELISA/CD4)
- IFN- γ , IL-2, TNF- α (Flow cytometry)
- CD8+ T cells (ELISPOT)
- Antibodies in lymphocyte supernatant (ALS)

Slides Credits:

Henry Boom

Chuck Daley

David Lewinsohn

Madhu Pai

PROTECT



them from

TUBERCULOSIS

Keep them away from sick people

Insist on plenty of rest

Train them in health habits

Consult the doctor regularly

This campaign made possible through the sale of Christmas Seals